



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Enhanced organic acid production in genetic engineered strains of *Aspergillus carbonarius*

Yang, Lei; Lübeck, Mette; Lübeck, Peter Stephensen

Publication date:
2014

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Yang, L., Lübeck, M., & Lübeck, P. S. (2014). *Enhanced organic acid production in genetic engineered strains of Aspergillus carbonarius*. Poster presented at 35th Symposium on Biotechnology for Fuels and Chemicals, Portland, Oregon, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Enhanced organic acid production in genetic engineered strains of *Aspergillus carbonarius*

Lei Yang, Mette Lübeck and Peter S. Lübeck

Section for Sustainable Biotechnology, Copenhagen Institute of Technology, Aalborg University, Denmark

Introduction

The filamentous fungus *Aspergillus carbonarius* is a good citric acid producer¹ and has great abilities in utilizing a broad range of substrates as well as excellent tolerance to stress conditions. It is selected for pathway engineering of the central carbon metabolism to improve the conversion of the renewable carbon sources and the flux into selected organic acids.

The present work focuses on development of two different mutants, a *gox* mutant and a *pepck* mutant. The *gox* mutant was constructed by deleting the *gox* gene which encode glucose oxidase in *A. carbonarius*. The *pepck* mutant was constructed by inserting the gene *pepck*, which encodes the phosphoenolpyruvate carboxykinase in *Actinobacillus succinogenes*, into *A. carbonarius*.

Objectives

- Deletion of the *gox* gene which is assumed to encode glucose oxidase in *A. carbonarius*
- Insertion of the gene *pepck*, which encodes the phosphoenolpyruvate carboxykinase in *Actinobacillus succinogenes*, into *A. carbonarius*
- Evaluation of the effect of two genetic modifications on organic acid production in fermentation



Method and Materials

1) Strains:

A. carbonarius wild-type 5010 was used to construct the *pepck* mutant and a Ku deficient and uracil auxotrophic transformant of the wild type strain, was used for construction of the *gox* mutant

2) Plasmids

Two plasmids pSB₁pepck and pSB₁gox built by USER cloning were used to construct *pepck* and *gox* mutant respectively.



3) Fermentation:

- Batch fermentation was carried out with the selected mutant in shaking flasks for 7 days by employing *A. carbonarius* wt 5010 and the Ku deficient strain (for comparison with the *gox* mutant only) as control.
- The acid production medium was as described by Goldberg² but supplemented with 0.1 g/L Zn.
- The concentration of gluconic acid and malic acid was analyzed by enzyme assays (Megazyme). The amount of glucose and other organic acids were analyzed by HPLC.

Results and Conclusion:

- a) As shown in Figure 1 and 2, after deleting the *gox* gene, *A. carbonarius* is incapable of accumulating gluconic acid in fermentation. Furthermore, the effect of the *gox* gene deletion on production of other organic acid was evaluated. An increase of malic acid and citric acid production was obtained as well as accumulation of oxalic acid after 7 days fermentation in the *gox* mutant.




Figure 1




Figure 2




Figure 3

- b) As shown in Figure 3, the two *pepck* mutants with highest expression level of the *pepck* gene showed improvement on production of malic acid and citric acid compared with wild type strain.

References:

¹Ghareib, M. 1987, "Assimilation of galacturonate by *Aspergillus carbonarius*", Folia Microbiologica 32: 211-215

²Battat, E., Peleg, Y., Bercovitz, A., Rokem, J.S. & Goldberg, I. 1991, "Optimization of L-malic acid production by *Aspergillus flavus* in a stirred fermentor", Biotech Bioeng 37: 1108-1116.